

ABSTRACT OF THE DISCLOSURE

Used as the positive electrode active substance of a lithium ion secondary cell is a mixture of a lithium-nickel-cobalt-manganese composite oxide represented by the formula

5 $\text{LiNi}_{(1-x-y)}\text{Co}_x\text{Mn}_y\text{O}_2$ wherein $0.5 < x+y < 1.0$ and $0.1 < y < 0.6$ and a lithium-manganese composite oxide represented by the formula $\text{Li}_{(1+z)}\text{Mn}_2\text{O}_4$ wherein $0 \leq z \leq 0.2$. The substance used gives outstanding power characteristics to the cell.

(21)
 (31)
 (41)
 (51)
 (61)
 (71)
 (81)
 (91)
 (101)
 (111)
 (121)
 (131)
 (141)
 (151)
 (161)
 (171)
 (181)
 (191)
 (201)
 (211)
 (221)
 (231)
 (241)
 (251)
 (261)
 (271)
 (281)
 (291)
 (301)
 (311)
 (321)
 (331)
 (341)
 (351)
 (361)
 (371)
 (381)
 (391)
 (401)
 (411)
 (421)
 (431)
 (441)
 (451)
 (461)
 (471)
 (481)
 (491)
 (501)
 (511)
 (521)
 (531)
 (541)
 (551)
 (561)
 (571)
 (581)
 (591)
 (601)
 (611)
 (621)
 (631)
 (641)
 (651)
 (661)
 (671)
 (681)
 (691)
 (701)
 (711)
 (721)
 (731)
 (741)
 (751)
 (761)
 (771)
 (781)
 (791)
 (801)
 (811)
 (821)
 (831)
 (841)
 (851)
 (861)
 (871)
 (881)
 (891)
 (901)
 (911)
 (921)
 (931)
 (941)
 (951)
 (961)
 (971)
 (981)
 (991)